

# *Molecular Dynamics Simulation of Water by TIP5P Model*

*Motohiko Tanaka, PhD., Nagoya, Japan*

*<http://www1.m4.mediacat.ne.jp/dphysique/>*

*Procedures of molecular dynamics simulation of water and ice are written for the five-point TIP5P model. Formerly, the shake/rattle method was used, showing the ice state not melted by microwaves – our theory discovery in J.Chem.Phys. 2007. Today, we use five-body water molecules with two hydrogens H1, H2 and two L1, L2 hydrogens of dummy sites. The fifth site of an oxygen site is used with Lennard–Jones potential  $\Psi = A/r^{12} - B/r^6$ . We regain similar results (values are some different), due to the structure of six-membered water and ice !*

1. “Classical Mechanics”, H. Goldstein, C. Poolee, J. Safko, 3rd Edition, Pearson Education Inc., England, 2003.
2. ”Microwave heating of water, ice and saline solution: Molecular dynamics study”, M.Tanaka and M.Sato, J.Chem.Phys., 126, 034509 1-9 (2007).

## *Procedures of Water Molecules by Five–Body Model*

- A. Five sites are oxygen( $O$ ), hydrogen 1 and 2( $H$ ), and hydrogen virtual  $L$  sites. They have, 0,  $+0.241e$ , and  $-0.241e$  charges, respectively. The  $L1$  and  $L2$  are called dummy sites.
- B. Separate  $\mathbf{R}_j$ ,  $\mathbf{V}_j$  and  $\mathbf{r}_i$  with  $j = 1, N / 5$ ,  $i = 1, N$  for molecules, and  $\mathbf{r}_i = (x_i, y_i, z_i)$  means for the three sites. The separation is done at the starting step only; once determined at  $t = 0$ , they become constant in time.
- C. The half time step is first executed for a predictor step, and the full time step is made for advance of time.
- D. Before the end of one step, the forces are calculated.  
The  $L$  sites are calculated by algebraic vector operation.
- E. After correction of quaternions, go to the beginning of the cycle.  
The leap – frog method is used for the plasmas and waters.

*Each step illustrates: translation (Step 1), rotation (Step 2–4), and adding the fields (Step 5–8).*

0. *Read positions  $(x, y, z), i = 1, N$ , and quaternions from the file, 'read(30)e0,e1,e2,e3',  $j = 1, N / 5$  (by Dr.Matsumoto, Okayama Univ).*
1. *Sum up five sites and advance,*  $\frac{d\mathbf{V}_j}{dt} = \frac{1}{m_j} \sum_{k=1}^5 \mathbf{F}_k$ ,  $\frac{d\mathbf{R}_j}{dt} = \mathbf{V}_j$ ,  
*for the translational motion.*
2.  $\frac{d\mathbf{L}_j}{dt} = \sum_{k=1}^5 (y_k F_k^z - z_k F_k^y, z_k F_k^x - x_k F_k^z, x_k F_k^y - y_k F_k^x)$   
*for the rotational motion : the sums are made over the five sites.*
3.  $\omega_{j,\alpha} = (A_{\alpha 1} L_x + A_{\alpha 2} L_y + A_{\alpha 3} L_z) / I m_{j,\alpha}$ , *the angular frequency for species  $A_{\alpha\beta}$  and inertia moments  $I m_{j,\alpha}$  at  $\alpha = x, y, z$ .*
4.  $\frac{d\mathbf{q}_j}{dt} = \frac{\Delta t}{2} (-e_1 \omega_x - e_2 \omega_y - e_3 \omega_z, e_0 \omega_x - e_3 \omega_y + e_2 \omega_z, e_3 \omega_x + e_0 \omega_y - e_1 \omega_z, -e_2 \omega_x + e_1 \omega_y + e_0 \omega_z)$ .

*(continued)*

5. Get a new rotational matrix  $A_{ij}(e_0, e_1, e_2, e_3)$  of the next time step below.

\*The predictor and corrector method is used in timings of time steps of Steps 2–5.

$$6. \mathbf{r}_i = \mathbf{R}_j + \begin{pmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{pmatrix} \begin{pmatrix} x_i \\ y_i \\ z_i \end{pmatrix} \text{ at the three sites } \mathbf{r}_i \text{ and the position } \mathbf{R}_j.$$

The dummy sites are calculated by algebraic operation.

7. Forces at Coulomb and LJ potentials are calculated using five sites.

8. Correction a normalization of quaternions is made at every 10 steps, and go to the next time step of Step 1.

Note that a time step is important. It will be  $\Delta t = 0.025$  or less, else the code is inaccurate or goes overflow.

# Quaternions in Place of Angles

$$e_0 = \cos \frac{\theta}{2} \cos \frac{\phi + \psi}{2}$$

$$e_1 = \sin \frac{\theta}{2} \cos \frac{\phi - \psi}{2}$$

$$e_2 = \sin \frac{\theta}{2} \sin \frac{\phi - \psi}{2}$$

$$e_3 = \cos \frac{\theta}{2} \sin \frac{\phi + \psi}{2}$$

*Classical Mechanics (3<sup>rd</sup> Edition)  
H. Goldstein , C.P. Poole, J.Safko,  
Pearson Education Inc., England 2003.*

**Only three of them are independent  
to avoid a gimbal lock**

Quaternion representation (4.47)

## Rotation matrix

$$\mathbf{r} = \mathbf{R} + \mathbf{A}^t \mathbf{r}'''$$

$$\mathbf{A} = \begin{pmatrix} e_0^2 + e_1^2 - e_2^2 - e_3^2 & 2(e_1 e_2 + e_0 e_3) & 2(e_1 e_3 - e_0 e_2) \\ 2(e_1 e_2 - e_0 e_3) & e_0^2 - e_1^2 + e_2^2 - e_3^2 & 2(e_2 e_3 + e_0 e_1) \\ 2(e_1 e_3 + e_0 e_2) & 2(e_2 e_3 - e_0 e_1) & e_0^2 - e_1^2 - e_2^2 + e_3^2 \end{pmatrix}$$

Time derivative of  
quaternions  
 $e_0, e_1, e_2, e_3$

$$\begin{pmatrix} \dot{e}_0 \\ \dot{e}_1 \\ \dot{e}_2 \\ \dot{e}_3 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -e_1 & -e_2 & -e_3 & e_0 \\ e_0 & -e_3 & e_2 & e_1 \\ e_3 & e_0 & -e_1 & e_2 \\ -e_2 & +e_1 & e_0 & e_3 \end{pmatrix} \begin{pmatrix} \omega_x \\ \omega_y \\ \omega_z \\ 0 \end{pmatrix}$$

# *The Lennard-Jones Potentials*

*With the Coulombic interactions, the 12–6 Lennard–Jones potential is adopted for the TIP4P and TIP5P cases:*

$$\Phi(r) = A / r^{12} - B / r^6$$

*for TIP4 :*

$$A = 4.17 \times 10^{-8} \text{ erg} \bullet \text{Ang}^{12}, B = 4.24 \times 10^{-11} \text{ erg} \bullet \text{Ang}^6$$

*for TIP5 – Ewald sum :*

$$A = 3.85 \times 10^{-8} \text{ erg} \bullet \text{Ang}^{12}, B = 4.36 \times 10^{-11} \text{ erg} \bullet \text{Ang}^6$$

*Some parameters are,*

$$r(OH) = 0.9572 \text{ Ang}, \Delta HOH = 104.52^\circ$$

$$r(OM) = 0.15 \text{ Ang for TIP4P only}$$

*The equipartition line of the virtual M site is on the plain that equally separates the HOH angle. The TIP5P cases are also available.*

## *To Start a Run*

*To start a simulation of water cluster with the TIP5P code, the adjacent 4x4 hydrogen pairs are summed electrostatically, and oxygen pairs are coupled by TIP5P Lennard-Jones potentials.*



*To get an initial state, we make the size of at least a 6x6x6 water cluster for numerical stability. Short-range and long-range Coulombic forces are best separated for interactions, and the short-range forces are made to be spatially dumped.*

*Around a given temperature, a dryrun is executed at least for 5 periods that is  $10^{-9}$  seconds. The long dryrun is very important !!*

*Then, we apply the electric field  $E_x = E_0 \sin(\omega * time)$  in the x-direction to excite the electric dipole interactions of water. For the moment, we give the electric field 10 GHz where the electric field  $E_0$  and electric dipole  $p_0$  are of the order of  $5 \times 10^{-3}$  eV.*

## To Obtain the Initial Equilibrium for 298 K

We will use salt ions of  $\text{Na}(+)$  and  $\text{Cl}(-)$  initially as the dryrun to give random noises. The  $6 \times 6 \times 6$  water clusters have 64  $\text{Na}(+)$  and 64  $\text{Cl}(-)$  ions, and a run time is  $t=1,700$ . We can see random water clusters. Afterwards, the salt ions are gradually removed, and the dryrun is continued for 5 periods up to  $t=50,000$ .



At start and the end of the dryrun of salt ions of  $t=0 - 3700$ . The dryrun without ions is continued afterward for 5 periods.

## **Check of quaternion: $A_{ij}$ coefficients**

$$\begin{aligned}
A_{11} &= (-\xi^2 + \eta^2 - \zeta^2 + \chi^2) = -\sin^2 \frac{\theta}{2} \sin^2 \frac{\psi - \phi}{2} + \sin^2 \frac{\theta}{2} \cos^2 \frac{\psi - \phi}{2} \\
&\quad - \cos^2 \frac{\theta}{2} \sin^2 \frac{\psi + \phi}{2} + \cos^2 \frac{\theta}{2} \cos^2 \frac{\psi + \phi}{2} \\
&= \sin^2 \frac{\theta}{2} \left( -\sin^2 \frac{\psi - \phi}{2} + \cos^2 \frac{\psi - \phi}{2} \right) - \cos^2 \frac{\theta}{2} \left( \sin^2 \frac{\psi + \phi}{2} - \cos^2 \frac{\psi + \phi}{2} \right) \\
&= \sin^2 \frac{\theta}{2} \left( 2 \cos^2 \frac{\psi - \phi}{2} - 1 \right) + \cos^2 \frac{\theta}{2} \left( \cos^2 \frac{\psi + \phi}{2} - 1 \right) \\
&= \sin^2 \frac{\theta}{2} (1 + \cos(\psi - \phi) - 1) + \cos^2 \frac{\theta}{2} (1 + \cos(\psi + \phi) - 1) \\
&= \sin^2 \frac{\theta}{2} \cos(\psi - \phi) + \cos^2 \frac{\theta}{2} \cos(\psi + \phi) \\
&= \sin^2 \frac{\theta}{2} (\cos \psi \cos \phi + \sin \psi \sin \phi) + \cos^2 \frac{\theta}{2} (\cos \psi \cos \phi - \sin \psi \sin \phi) \\
&= \frac{1}{2} (1 - \cos \theta) (\cos \psi \cos \phi + \sin \psi \sin \phi) + \frac{1}{2} (1 + \cos \theta) (\cos \psi \cos \phi - \sin \psi \sin \phi) \\
&= \cos \psi \cos \phi - \cos \theta \sin \psi \sin \phi
\end{aligned}$$

$$\begin{aligned}
A_{12} &= 2(\xi\chi - \xi\eta) = 2(\cos^2 \frac{\theta}{2} \sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2} - \sin^2 \frac{\theta}{2} \sin \frac{\psi - \phi}{2} \cos \frac{\psi - \phi}{2}) \\
&= (1 + \cos \theta) \sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2} - (1 - \cos \theta) \sin \frac{\psi - \phi}{2} \cos \frac{\psi - \phi}{2} \\
&= \sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2} - \sin \frac{\psi - \phi}{2} \cos \frac{\psi - \phi}{2} + \cos \theta (\sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2} \\
&\quad + \sin \frac{\psi - \phi}{2} \cos \frac{\psi - \phi}{2}) \\
&= \frac{1}{2} \sin \left( \frac{\psi + \phi}{2} + \frac{\psi + \phi}{2} \right) - \frac{1}{2} \sin \left( \frac{\psi - \phi}{2} + \frac{\psi - \phi}{2} \right) + \frac{1}{2} \cos \theta (\sin(\psi + \phi) \\
&\quad + \sin(\psi - \phi)) \\
&= \frac{1}{2} \sin(\psi + \phi) - \frac{1}{2} \sin(\psi - \phi) + \frac{1}{2} \cos \theta (\sin(\psi + \phi) - \frac{1}{2} \sin(\psi - \phi)) \\
&= \frac{1}{2} (\sin \psi \cos \phi + \cos \psi \sin \phi - \sin \psi \cos \phi + \cos \psi \sin \phi) + \frac{1}{2} \cos \theta (same) \\
&= \cos \psi \sin \phi + \cos \theta \sin \psi \cos \phi
\end{aligned}$$

$$\begin{aligned}
A_{13} &= 2(\eta\zeta + \xi\chi) = 2\left(\sin\frac{\theta}{2}\cos\frac{\theta}{2}\cos\frac{\psi-\phi}{2}\sin\frac{\psi+\phi}{2}\right. \\
&\quad \left. + \sin\frac{\theta}{2}\cos\frac{\theta}{2}\sin\frac{\psi-\phi}{2}\cos\frac{\psi+\phi}{2}\right) \\
&= 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\left[\cos\frac{\psi-\phi}{2}\sin\frac{\psi+\phi}{2} + \sin\frac{\psi-\phi}{2}\cos\frac{\psi+\phi}{2}\right] \\
&= \sin\theta\left[\sin\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2} + \cos\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right] \\
&= \sin\theta\left[\sin\left(\frac{\psi+\phi}{2} + \frac{\psi-\phi}{2}\right)\right] \\
&= \sin\theta\sin\psi
\end{aligned}$$

$$\begin{aligned}
A_{21} &= -2(\xi\eta + \zeta\chi) = -2(\sin^2 \frac{\theta}{2} \cos \frac{\psi - \phi}{2} \sin \frac{\psi - \phi}{2} \\
&\quad + \cos^2 \frac{\theta}{2} \sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2}) \\
&= -(1 - \cos \theta) \cos \frac{\psi - \phi}{2} \sin \frac{\psi - \phi}{2} - (1 + \cos \theta) \sin \frac{\psi + \phi}{2} \cos \frac{\psi + \phi}{2} \\
&= -\frac{1}{2}(1 - \cos \theta) \sin(\psi - \phi) - \frac{1}{2}(1 + \cos \theta) \sin(\psi + \phi) \\
&= -\frac{1}{2}(\sin(\psi - \phi) + \sin(\psi + \phi)) + \frac{1}{2}\cos \theta (\sin(\psi - \phi) - \sin(\psi + \phi)) \\
&= -\frac{1}{2}(sc - cs + sc + cs) + \frac{1}{2}\cos \theta (sc - cs - sc - cs) \\
&= -\sin \psi \cos \phi - \cos \theta \cos \psi \sin \phi
\end{aligned}$$

$$\begin{aligned}
A_{22} &= \xi^2 - \eta^2 - \zeta^2 + \chi^2 \\
&= \sin^2 \frac{\theta}{2} (\sin^2 \frac{\psi - \phi}{2} - \cos^2 \frac{\psi - \phi}{2}) + \cos^2 \frac{\theta}{2} (-\sin^2 \frac{\psi + \phi}{2} + \cos^2 \frac{\psi + \phi}{2}) \\
&= \sin^2 \frac{\theta}{2} (1 - 2 \cos^2 \frac{\psi - \phi}{2}) + \cos^2 \frac{\theta}{2} (2 \cos^2 \frac{\psi + \phi}{2} - 1) \\
&= -\sin^2 \frac{\theta}{2} \cos(\psi - \phi) + \cos^2 \frac{\theta}{2} \cos(\psi + \phi) \\
&= -\frac{1}{2} (1 - \cos \theta) \cos(\psi - \phi) + \frac{1}{2} (1 + \cos \theta) \cos(\psi + \phi) \\
&= \frac{1}{2} (-\cos(\psi - \phi) + \cos(\psi + \phi)) + \frac{1}{2} \cos \theta (\cos(\psi - \phi) + \cos(\psi + \phi)) \\
&= \frac{1}{2} (-cc - ss + cc - ss) + \frac{1}{2} \cos \theta (cc + ss + cc - ss) \\
&= -\sin \psi \sin \phi + \cos \theta \cos \psi \cos \phi
\end{aligned}$$

$$\begin{aligned}
A_{23} &= 2(\eta\chi - \xi\zeta) = 2\left(\sin\frac{\theta}{2}\cos\frac{\theta}{2}\cos\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2}\right. \\
&\quad \left.- \sin\frac{\theta}{2}\cos\frac{\theta}{2}\sin\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right) \\
&= 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\left(\cos\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2} - \sin\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right) \\
&= 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\cos\left(\frac{\psi+\phi}{2} + \frac{\psi-\phi}{2}\right) \\
&= \sin\theta\cos\psi
\end{aligned}$$

$$\begin{aligned}
A_{31} &= 2(\eta\zeta - \xi\chi) = 2\left(\sin\frac{\theta}{2}\cos\frac{\theta}{2}\sin\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2}\right. \\
&\quad \left.- \sin\frac{\theta}{2}\cos\frac{\theta}{2}\cos\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right) \\
&= 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\left(\sin\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2} - \cos\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right) \\
&= 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\sin\left(\frac{\psi+\phi}{2} - \frac{\psi-\phi}{2}\right) \\
&= \sin\theta\sin\phi
\end{aligned}$$

$$\begin{aligned}
A_{32} &= -2(\xi\zeta + \eta\chi) = -2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\left(\sin\frac{\psi+\phi}{2}\sin\frac{\psi-\phi}{2}\right. \\
&\quad \left.+ \cos\frac{\psi+\phi}{2}\cos\frac{\psi-\phi}{2}\right) \\
&= -\sin\theta\cos\left(\frac{\psi+\phi}{2} - \frac{\psi-\phi}{2}\right) \\
&= -\sin\theta\cos\phi
\end{aligned}$$

$$\begin{aligned}
A_{33} &= -\xi^2 - \eta^2 + \zeta^2 + \chi^2 \\
&= -\sin^2 \frac{\theta}{2} \left\{ \sin^2(-) + \cos^2(-) \right\} + \cos^2 \frac{\theta}{2} \left\{ \sin^2(+) + \cos^2(+) \right\} \\
&= -\sin^2 \frac{\theta}{2} + \cos^2 \frac{\theta}{2} = \frac{1}{2} \left\{ -(1 - \cos \theta) + 1 + \cos \theta \right\} = \cos \theta \quad QED!
\end{aligned}$$